



New diseases and health risk in cage free systems

Fernando Carrasquer Puyal
DVM CEAV ECPVS Resident
Global Technical Service – Veterinary specialist
H&N International GmbH

What makes the pathology in cage-free birds different?



Faecal-oral infection is possible



Some behaviours are not mitigated



Low biosecurity on external parks



Flock management is more complicated

Most important health problems in US

Caged laying hens


Rank	Disease	Score
1 tie	E coli	2.63
1 tie	Infectious bronchitis	2.63
3	Infectious coryza	2.37
4 tie	FDN	2.16
4 tie	ILT	2.16
4 tie	Calcium Depletion	2.16
7	Coccidiosis	1.89
8 tie	Cannibalism	1.79
8 tie	NFM	1.79
8 tie	Necrotic enteritis	1.79

Cage-free laying hens

Rank	Disease	Score
1	E coli	3.28
2 tie	Infectious bronchitis	2.78
2 tie	Roundworms	2.78
4 tie	Piling	2.72
4 tie	Cannibalism	2.72
6	Coccidiosis	2.33
7	ILT	2.28
8	Infectious coryza	2.17
9	Necrotic enteritis	2.00
10 tie	FDN / FC / EDS	1.89




Coccidiosis



Coccidiosis


Etiologic agent: **Eimeria Spp.**

- It is a protozoa that needs to cycle in the environment and in the poultry gut
- Different species produces different lesion in the gut
- It is present worldwide
- There is no cross protection between species



Coccidiosis

Eimeria Tenella




Coccidiosis

Eimeria maxima



Coccidiosis

Eimeria Acervulina



Coccidiosis

Eimeria Brunetti



Coccidiosis


Eimeria necatrix



Coccidiosis


Infectious form: **oocyst**

- Very resistant in the environment
- Heavy and big size
- It needs to sporulate to become infective in a humid and warm ambiance
- Different species differs in size and morphology



Coccidiosis

Gut health & Coccidia



Source:
Alnassan 2014

Coccidiosis

Health program objective

Short life birds



No delayed growth
No anticocci resistance



1. Anticocci program

Rotation program

Shuttle program

Bio-shuttle program

2. Vaccine programs

3. Phyto programs

Long life birds



Long lasting immunity
against the different
Eimeria species



1. Vaccines program

+ Anticocci

+ Phyto

2. ~~Anticocci programs~~

Challenge required !!!

Coccidiosis

Vaccine by type of bird



Short life birds vaccines

Eimeria acervulina
Eimeria maxima
Eimeria Tenella
Eimeria Mitis
...



Long life birds vaccines

Eimeria acervulina
Eimeria maxima
Eimeria tenella
Eimeria mitis
Eimeria brunetti
Eimeria praecox
Eimeria Necratix

Coccidiosis

Vaccine by strain attenuation



Natural strains

- +** - Low cost
- +** - Excellent dissemination in the field
- - Still pathogenic strains: mitigation may be required
- - Introduction of pathogenic strains into the house




Attenuated strains

- +** - Less lesions produced during the Eimeria cycle
- +** - Non pathogenic strains
- - High cost
- - Poor dissemination in the field

Coccidiosis


Attenuated vaccines VS Natutal vaccines



Source:
M. Dardi

Coccidiosis

Attenuation by precocity method



Coccidiosis

Vaccine administration methods



~~Spray in feed~~



Spray chick boxes
at farm



~~Drinking water
(Nipples)~~



Spray chick boxes
at hatchery

Coccidiosis

Spray vaccine process



Vaccine preparation



Coarse Spray



Droplet ingestion



Vaccine strain recirculation

Coccidiosis

Coarse spray: The size (and the weight) matters




Coccidiosis

Droplet ingestion: using the curiosity to our advantage

What is this red thing in your neck?


Whoaa, red things !!!



Beautiful droplet
Curious chicks
Light
Time
Temperature

Coccidiosis

Vaccine recirculation



Coccidiosis

Vaccine recirculation




Natural strains


- **Assure strain dissemination**
- **Mitigate strains pathogenicity**
 - Give more room ASAP
 - Control litter humidity
 - **LAST RESOURCE:**
Anticocci treatment



Attenuated strains

- **Facilitate strain dissemination**
 - Give more room while observing the Eimeria cycles.
 - When giving more room, move litter with the chicks
 - Assure litter humidity litter





Helminths

(Gastrointestinal parasite)

Helminths

What are we talking about?




Nematodes
Round worms



Pic :Poultry site

Cestodes
Tape worms




Pic :Animal parasitosis

Trematodes
Flukes

Helminths

Prevalence by production type – Systematic review




Source:
Shifaw 2021

Helminths

Nematodes (Round worms)


- Ascaris
 - Ascaridia sp.
 - Heterakis sp.
- Capillaires
 - Capillaria sp.
- Spirures
- Strongles
 - Trichostrongylus tenuis



Picture:
Reussir.fr

Ascaris

Ascaris species prevalence in European organic farms



Ascaris

Ascaridia

Etiologic agent:

Ascaridia Galli.

- Nematodes. It can measure 6—11 cm as adult and infest the intestine
- Egg drop and bodyweight losses is possible in case of strong infestation
- No report of infestation in humans
- It can be hosted by earth worms



Pictures:
Wattagen

Ascaris

Clinical sign


Anorexia
Loss of body weight
Ruffled feathers
Drooped wings
Retarded growth,
Altered hormone levels,
Depression,
Increased cannibalism

Not bad
for a
worm!!



Ascaridia

Effect of different infestation levels of *Ascaridia galli* on egg mass.



Source:
Sharma 2017

Ascaris

Heterakis

Etiologic agent:

Heterakis gallinarum

- Nematode. It can measure 1-1,5 cm and infest ceca.
- It can produce ceca inflammation
- It can host *Histomonas meleagridis*



Pictures:
Wattagen

(Beyond) Ascaris

Histomonosis

Etiologic agent:

Histomonas meleagridis.

- Flagellated amoeboid Protozoan
- Sulfur-colored droppings, characteristic lesion in ceca and liver
- High mortality (30%) can occur in chickens
- Very complicated treatment because the lack of authorized drugs



Ascaris

Health program

Prevention

Biosecurity

(Free range???)

Eradication

Not possible

Control

Monitoring

Ascaris finding in autopsies

Egg in faecal droppings

~~Vaccines~~

Treatment

Flubendazol

Piperazine

...

Passive control

Specific disinfection

Outdoor park rotation ?

Brachyspira

Brachyspira

Etiologic agent:

Brachyspira Piloscoli

Brachyspira intermedia

- Reduced egg production, downgrading of shell eggs, bodyweight loss
- Most common in free range birds
- Foamy yellowish ceca content



Brachyspira


Prevalence by production type –



Source:
Hess 2017

Brachyspira

Raw materials effect



Source:
Phillips 2012

Brachyspira

Health program

Prevention

Biosecurity

(Free range???)

Eradication

Not possible

Control

Avoid colonization

- C&D protocol
- Strict biosecurity routines
- Rodent control

No vaccines are currently available

Antimicrobial treatment.

- Tiamuline
- Tetraciline ?




Piling




Piling

Nest Piling



Piling

Panic piling



Predation episode



Loud sounds



Disturbing visits

Piling

Reiterative piling

First pile-up event



Pile-up increase event



Piling event



Suffocation event



Source:
Gray 2020

Bonus track Toe-Pecking

What is it about?


- Only in white layers
- Different Breeds
- No beak treatment
- Mortality from 0,1%-0,3%



Netherlands Germany Switzerland Finland



Thank you for your attention



H&N International – Making your success the center of our universe



Follow us on LinkedIn
H&N International GmbH



Find out more about
KAI farming assistant