



Cage-Free Rearing

H&N Cage-Free academy 2023

Leon Schouren
Key Account Manager Europe & Global Technical Service
schouren@hn-int.com

Production systems Global



Cages




Enriched cages



Barn egg



Aviary systems



Free range



Biological

Cage-Free Rearing

Influences on Layer Behavior



Lighting



Feed



Red mites



Vaccination



Climate



Flock density

Cage-Free Rearing

What do we like to know before we start rearing?

- White or Brown layers?
- Production based on number of eggs, Kg of eggs, market needs
- Age of the Layers in production
- Type of equipment rearing/production
- Type of feeding system rearing/production
- Daylight influence rearing/production



Cage-Free Rearing

Make a plan before the start of rearing

Determine the strategic road with:


Advisors from the hatchery, feed suppliers, Veterinarian and other involved parties!




EQUIPMENT REQUIREMENT

Aviary systems with feed / water / nest boxes on different levels

- With this type of system, the layers must jump on different levels to find feed / water / nest boxes.
- We prefer to train all the pullets to learn how to move in this type of production system in rearing period!

-  **The best kind of rearing you need for this system**
- Aviary systems within height adjustable slats
 - Aviary row systems
 - Aviary row systems within height adjustable slat inside the system



WHAT KIND OF REARING SYSTEM & TRAINING PROGRAM YOU NEED

Try to plan ahead to use the best type of rearing system for the layer house.

H&N Cage-Free
Management Guide
<https://hn-int.com/>

Equipment Requirement

Cage-Free Rearing

- The more closely the rearing facility resembles the future production system, the easier it will be for the pullets to settle down in their new environment after being transferred to the laying house.
- With this we can use the complete genetic potential of our H&N breed.



Equipment Requirement

Cage-Free Rearing

- Barn system with full litter
- Barn system with full slats
- Barn system with 2/3 slats and 1/3 litter
- Aviary systems with in height adjustable slats
- Aviary row systems
- Aviary row systems with in height adjustable slat(s) inside the system.

EQUIPMENT REQUIREMENT



STOCKING DENSITY

An adequate stocking density leads to success in rearing chicks. A high stocking density impacts negatively on daily growth, flock uni-

formity and chick development. Furthermore, a high stocking density combined with reduced feeder space will limit feed consump-

tion, which might already be low under certain conditions (e.g. hot climate or poor feed quality) and sufficient access to water.

Table 2: Stocking density in rearing farms*

Age	Floor space		Feeder space**		Drinker Space	
	Row-systems	Floor***	Row-systems	Floor	Row-systems	Floor
0 – 3 weeks	140 cm ² /bird	21 birds/m ²	2.5 cm/bird	4 cm/bird 60 birds/pan	1.25 trough cm/bird 12 birds/nipple	1.4 trough cm/bird 12 birds/nipple 100 birds/fountain
3 – 16 weeks	285 cm ² /bird	16 birds/m ²	5 cm/bird	8 cm/bird 30 birds/pan	2.5 trough cm/bird 8 birds/nipple	2.5 trough cm/bird 8 birds/nipple 75 birds/fountain

*This table is a general recommendation and you should adhere to your own country's recommendations.

**minimal recommendations


*** includes all the available space

**H&N Cage-Free
Management Guide**
<https://hn-int.com/>

Aviary Rearing

In Height adjustable systems

- ✗ Planning white/brown pullets
- ✗ Catching pullets
- ✗ Mortality by accidents
- ✗ No manure belt
(not used to sound)
- ✗ Higher costs/hen



Aviary Rearing

Row systems

- ✗ Learn how to jump 100%!!!!
- ✗ Enrichment first days in cages
- ✗ Ventilation block by system



Aviary Rearing!



EQUIPMENT REQUIREMENT (7 STEPS)

1. What kind of feeding, and drinking system is used in production?
2. Do the layers need to move on slats in production?
3. Is feed/water/nest boxes placed on the same level in production?
4. Do the layers need to jump in production system to find feed/water/nestboxes on different levels?
 1. Do the birds need to jump on perches to find these different levels?
 2. What is the maximum height that the layers need to jump to in the production system?
 3. Are there manure belts in the production system?

**H&N Cage-Free
Management Guide**
<https://hn-int.com/>

Vaccination Program



KAI



Discuss with local veterinarian for best choice.
Also looking at economic benefits!
**Try to finish vaccination program before
transfer!**

Brooding

Prepare for housing

- Preparing for placement from DOC
 - *Check setting in the computers if they are O.K.*
 - *Is everything checked & tested (especially with first flock)*
- Start pre-heating the house on time.
 - *House temperature up to 35°-36° C.
In summer time, at least 24 hours before arrival of the chicks.
In wintertime **at least** 48 hours before arrival of the birds*
 - *The first 48 – 72 hours after placement, this temperature must be maintained.*

The relative humidity preferred to be at least 60%.




Brooding

temperature




KAI



Brooding

temperature




Measuring Chick Body Temperature
Optimum 40-41° C (104-106° F)

Adjust House Temperature according to the Chick Body Temperature!

Brooding

- Check the drinker height regularly, especially in first weeks of Rearing
- Give extra attention with IR treatment!
360° nipple drinker/cup drinkers



Too Low



Too High



Right Height

Brooding


Intermittent lighting program

Intermittent Lighting Program

► Dark houses only (< 3 lux)


This program can be used for up to 7 – 10 days after arrival. Then switch back to the regular step-down lighting program. Using this lighting program has the following advantages:

- Chick behaviour is synchronized; they rest or sleep at the same time.
- Weak chicks will be stimulated by stronger ones to move as well as to eat and drink.
- The behaviour of the flock is more uniform and evaluating the flock is much easier.
- Chick mortality will decrease.



Lighting Programs

Should this program be implemented for ALL the flock worldwide?





**The same lighting program
cannot be used worldwide**

1. Effects of the natural light
2. Possibility to adapt egg weight to different market demands by using a targeted lighting program

Lighting Program

influencing factors



- Latitude/ Longitude: *North/South, East/West*
- Hatch date: *Year Season, Natural day light*
- House design: *Closed, light tight, open sided*
- Target body weight: *Egg size, feed consumption*



Lighting Programs

Comparing a Slow Down versus a Fast Down Program

If it doesn't appear that body weight targets will be met at week 5, it is highly recommended to change to a slower reduction to allow body weight to improve!



Lighting Program

Step Down Period

- Step wise reduction of day length from placement until age of 4 to 8 weeks.
- Around 3-10 weeks, be extra aware of pecking, then reduce the light intensity. (Temporarily)
- The day length will remain constant on 8-12 hours
- Don't stick to the lighting schedule. If the birds don't reach their bodyweight, don't reduce the light further until they are back of target



Lighting Program

Constant Period

- The day length during the Constant Period should not be chosen too short. (8-12 hours)
- The birds still have to grow.
They need some time to eat to supply bodyweight gain.
- Never increase the day length during this period.
This can induce an advanced onset of lay.



Lighting Program

Measure number of LUX, avoid bright light spots (feather pecking) Min 15 LUX in rearing for brown pullets and 10 LUX for white pullets!!!!




Development of the chicks

The birds (and not only ours!) have one failure: They can't read!

- *No management guide*
- *No lighting program*

Therefore, it is the pulletgrower who has to determine if everything is „on track“



- *Take bodyweights from 1st week onwards, once a week*
- *Not necessary to take individual BW, only if necessary.*

Development of the chicks

- Start training with lighting (Dimming period) as soon as possible after the intermittent lighting program. (14 days)
- Let the chicks out of the system a.s.a.p!!!!
- Be aware of situations with paracocc vaccinations, recycled paper!!



Development of the chicks




Development of the chicks

- At each change, check on the behaviour of the pullets and how they react to the change.

(e.g., Open the system, change feeding times or increasing the height of the nipple drinkers!

- Very important: make sure that 100% of the pullets are active!
- It is very important that **ALL** the pullet move upwards to the slats/system and perches.

Body Development During Rearing



FEED MANAGEMENT

- Target to feed ALL the birds the same quantity, and in this quantity the COMPLETE balanced feed.
- To prepare the birds for the big step in feed intake at the start of production.
- To keep up feeding in hot climate areas
- To reduce select feed intake
- To prevent restrict feeding



**H&N Cage-Free
T.T. Feed Management**
<https://hn-int.com/>

New challenges in cage-free

Does it work in ps/commercial rearing/layer farms, long feed chains & high stocking densities ?



New challenges

Cage versus Cage-Free



New challenges

Ban on beaktreatments



IR-treatment



Untrimmed

Feed Structure

Good



**Homogenous
mash feed structure
the basis
for good & even feed
and nutrient intake**

Feed structure

Judged by a hen

- ☹️ I don't like hard and sharp granulate
- ☹️ I don't like fine powder mashfeed
- ☹️ I don't like hard pallets ...



Source: G. G. Mateos

Feed structure

What are they going to eat?



Feed structure

Easy testing on farm level



Feed structure

Easy testing on farm level



Invest into the right feeding system

And take the profit!

- feed chain
 - Fast running feed chain (>18 M/Min)
 - Enough capacity for transport augers or feed hoppers
 - flat bottom silos!
-
- ✓ **Profit from less to no waste**
 - ✓ **Better feeding behaviour**




Feed Management

When to start feed management?

- In rearing period (age of 3-4 weeks)
- Start to empty the feeders first one or two times a week. (Keep an eye on behaviour of the flock)
- From 6 weeks of age, we let them eat the feeders till low level, **ones every day.**
- **Continue this also after transfer to production!**



FEED MANAGEMENT



Rearing

Alfalfa (Luzerne)

Use the see if everything running smoothly!



Alfalfa (Luzerne)

Use the see if everything running smoothly!



- Do we need to use Alfalfa?
- For what purpose do we need to use Alfalfa?
- When do we need to use Alfalfa?

Alfalfa (Luzerne)

Use the see if everything running smoothly!

- We don't use alfalfa to feed our birds!
- We use alfalfa only to give us some extra hints to see if everything is in good balance!
- **When birds don't take alfalfa, they are in good balance.**
- Advisable to use it during most stressfull periods:
 - In rearing 3 till 10 weeks



Feed Management

Mandatory Steps

- **From week 4** in rearing start to train the chicks to empty to feeding troughs 2-3 times a week!
- **From week 5-6** empty (low level) the feeding troughs every day!
- Continue this training after transfer in production!



(Cage-Free) Rearing

Summarized

- Good feed and feed management
- Observe the birds and they will “tell you”
- Rearing is the key to success




Please be excellent because ...

*... an excellent manager will still be able
to perform with acceptable results even when
circumstances are not perfect*

But ...

*A manager with poor skills will
spoil the birds and the production results
even in the best circumstances!*

Thank you for your attention



H&N International – Making your success the center of our universe



Follow us on LinkedIn
H&N International GmbH



KAI



KAI

Find out more about
KAI farming assistants

